



Quality Improvement

Quality Improvement Study Using a Machine Learning Mortality Risk Prediction Model Notification System on Advance Care Planning in High-Risk Patients

Jonathan Walter, MD¹, Jessica Ma, MD¹, Alyssa Platt², Yvonne Acker³, Mark Sendak⁴, Michael Gao⁴, Matt Gardner⁴, Suresh Balu⁴, Noppon Setji, MD¹

¹ Department of Medicine, Duke University School of Medicine,

² Department of Biostatistics, Duke University,

³ Duke University Health System,

⁴ Duke Institute for Health Innovation

Journal of Brown Hospital Medicine

Article Information

Keywords: Quality improvement, Machine learning model, Advance care planning, ACP

<https://doi.org/10.56305/001c.120907>

Submitted: January 19, 2024 EDT

Accepted: July 03, 2024 EDT

Abstract

Background: Advance care planning (ACP) is an important aspect of patient care that is underutilized. Machine learning (ML) models can help identify patients appropriate for ACP. The objective was to evaluate the impact of using provider notifications based on a ML model on the rate of ACP documentation and patient outcomes.

Methods: This was a pre-post QI intervention study at a tertiary academic hospital. Adult patients admitted to general medicine teams identified to be at elevated risk of mortality using a ML model were included in the study. The intervention consisted of notifying a provider by email and page for a patient identified by the ML model.

Results: A total of 479 encounters were analyzed of which 282 encounters with a notification sent to a provider met inclusion criteria. The covariate-adjusted proportion of higher-risk patients with documented ACP rose from 6.0% at baseline to 56.5% (Risk Ratio (RR)= 9.42, 95% CI: 4.90 - 18.11). Patients with ACP were more than twice as likely to have code status reduced when ACP was documented (29.0% vs. 10.8% RR=2.69, 95% CI: 1.64 - 4.27). Additionally, patients with ACP had twice the odds of hospice referral (22.2% vs. 12.6% Odds Ratio=2.16, 95% CI: 1.16 - 4.01). However, patients with ACP documented had a longer mean LOS (9.7 vs. 7.6 days, Event time ratio = 1.29, 95% CI: 1.10 - 1.53)

Conclusion: Provider notifications using a ML model can lead to an increase in completion of ACP documentation by frontline clinicians in the inpatient setting.

BACKGROUND

Advance care planning (ACP) is an integral component of patient care. ACP emphasizes a patient's values, goals, and preferences regarding their medical treatment.¹ It has been shown to increase patient quality of life, decrease aggressive care at the end-of-life, and increase hospice utilization.²⁻⁶ Despite evidence supporting ACP for goal-concordant care, ACP is underutilized and often not initiated, especially in hospitalized patients.^{7,8} Further, documentation of patient wishes related to ACP in the electronic health record (EHR) is inconsistent, sparse, and difficult to find.⁹ This may be due to limited time and resources available to front-line physicians to initiate ACP and document these discussions.^{10,11}

Another barrier to appropriate initiation of ACP is prognostication because providers often overestimate longevity and are too optimistic of patient prognoses.^{12,13} These inaccurate predictions can lead to delayed or

missed end-of-life conversations. To improve prognostication, machine learning (ML) models can support the identification of patients who may benefit from ACP and nudge providers to complete and document ACP conversations.¹⁴⁻¹⁶ Identification of these patients by ML models can also assist with appropriate timing and initiation of ACP. ML mortality risk prediction models have been integrated into outpatient cancer clinics and palliative care consultation services to improve ACP documentation, however limited data is investigating ACP conversations by hospitalist providers.^{17,18}

At our institution, we found that less than 5% of patients in 2017-18 had a documented ACP conversation in the EHR within the last six months of life. To improve ACP documentation, we developed a quality improvement (QI) intervention using an ML mortality risk prediction model. Based on the model's output, general medicine inpatient providers were notified to consider ACP when patients at elevated risk of mortality were ad-

mitted.¹⁹ In this paper, we describe the intervention and evaluate documentation of ACP conversations in high-risk patients.

METHODS

The primary aim of this study was to evaluate whether a notification based on an ML mortality model increases ACP documentation. Secondary aims sought to evaluate whether documentation of ACP was associated with differences in patient outcomes, including length of stay (LOS), 30-day readmission, intensive care unit (ICU) admissions, change in code status (new Do Not Attempt Resuscitation order (DNAR)), and discharge to hospice.

Study Design and Setting

This is a pre-post QI study at Duke University Hospital (DUH), a tertiary academic medical center, starting from January 2019 through April 2021. Patients were included if they were at least 18 years of age at admission, admitted to the general medicine service at DUH, and were deemed to have an elevated risk of mortality within 30 days of admission by a validated risk measurement tool developed by the Duke Institute for Health Innovation.¹⁹

The study included an intervention pilot phase from November 2019 through February 2020 that was initiated on general medicine hospitalist teams. In March 2020, the study was expanded to general medicine teaching service lines that included residents. The ML model identified patients appropriate for ACP by predicting their risk of mortality. Patients were classified into the categories of “low”, “medium”, “high”, or “critical” for mortality at the time of admission (Supplemental Table 1).²⁰

Once a patient was identified as medium or higher risk, the patient was screened for exclusion criteria, listed below, by the QI team administrator reviewing the model dashboard. Providers were then notified via text page and a templated email that they were caring for a patient that may benefit from ACP. After March 2020, the population was refined to general medicine patients with a 30-day mortality risk classification of “medium,” “high,” or “critical,” and a 6-month mortality risk classification of “high” or “critical” and continued until April 2021.

In addition to the notification, the intervention directed providers to use a dedicated ACP note template.²¹ The template included prompts to assess patient illness understanding, explore patient goals, fears, and worries, address, and document code status, and identify a surrogate decision maker (Appendix 1). The template was updated in January 2021 based on direct provider feedback, and these additions are noted in the appendix. Hospitalists received optional ACP education during a faculty meeting, and educational materials were included in each notification email. Ancillary teams were also included in

the notification to help improve the care for these high-risk patients. This included the pharmacy, case management, and documentation and coding teams.

Providers were eligible for notification if they were internal medicine residents, advance practice providers, or attending physicians at DUH. Patients were excluded if they were admitted to the ICU within 24 hours of admission, to observation status, or to a non-general medicine service. In the post period, patients were also excluded if they had a recently completed ACP note, had an established comfort care directive, the patient was discharged or planned for discharge prior to screening by the QI team administrator, or due to technological issues such as server downtime or model maintenance.

A closed and open-ended questionnaire was sent in August 2020 to hospital medicine providers to obtain provider feedback regarding the ACP notification intervention. Providers were asked to rate (strongly agree, agree, disagree, strongly disagree) the ACP notification’s impact on patient care, the accuracy of identified patients, and their satisfaction with the notification. A reminder to complete the survey was sent twice at two-week intervals.

The Duke University Institutional Review Board (IRB) approved and determined this study exempt (IRB Pro00104527).

Study Outcomes and Measures

The primary outcome was documentation of ACP conversations with at-risk patients. Providers were asked to document their conversation using a developed template, which was then queried to ascertain whether the primary outcome occurred (binary yes/no).

Secondary outcomes included LOS in days from inpatient admission to discharge alive. Change in code status was a binary indicator that code status changed from more intervention to less intervention during the hospital stay indicated by a new DNAR order from the time of admission to the time of discharge. ICU transfer was a binary indicator that a patient was transferred to or received ICU level of care during their stay. Since hospice referral was contingent on being discharged alive from the hospital, the outcome of hospice referral was defined as a 3-category nominal variable with levels defined as inpatient death, discharge without hospice referral, and discharge to hospice. Readmissions within 30-days of initial discharge were defined following the Centers for Medicare Services Hospital-Wide All-Cause readmission definition for unplanned readmissions to our health system and quantified for analysis by the length of time (in days) from hospital discharge alive to unplanned readmission to the same facility. Patient demographic information and clinical details were extracted from the EHR.

COHORTS

A historical control cohort was created by identifying patients discharged from general internal medicine hospitalist providers at DUH from January 1st 2019 until October 31st 2019 using the same machine-learning model. The identified patients were those identified using the mortality risk threshold classifications of “medium”, “high”, or “critical” at 30-days and “high” or “critical” at 6-months (supplemental Table 1).²⁰

We analyzed the primary outcome during the pre- and post-intervention periods and in 5 distinct time periods: pre-intervention (1/1/2019-11/17/2019), pilot phase (11/18/2019-2/14/2020), start of notifications to teaching teams (3/26/2020-6/25/2020), pause in teaching team notification for new resident education (6/26/2020-7/21/2020), and teaching team notification resumes (7/21/2020-4/30/2021) (Supplemental figure 1).

Statistical Analysis

The primary analysis examined the probability of ACP documentation before and after intervention implementation, using 5 distinct intervals described above. Probability of ACP documentation was modeled using Modified Poisson regression to approximate risk ratios (RR), adjusting for patient- and faculty-level confounders, using generalized estimating equations (GEE) with exchangeable working correlation and sandwich standard errors clustered at the provider level.²² Estimated RRs measure change in probability of receiving ACP from the pre-period to each follow-up period. Risk differences were derived using the delta method. An unadjusted intraclass correlation coefficient (ICC) for provider was calculated using the one-way ANOVA method.²³

Secondary analysis examined associations between ACP documentation and patient outcomes. Propensity score overlap weighting methods were used to balance patient characteristics between those that did and did not receive ACP, with mixed effects logistic regressions to predict receipt of ACP, and random intercepts at the level of the provider.²⁴ Standardized differences were computed to compare characteristics of patients before and after weighting, with values smaller than ± 0.10 considered acceptable.²⁵

In this high-risk population, death may be considered a semi-competing risk with some outcomes; therefore, LOS and readmissions were analyzed using survival analysis methods. For LOS, accelerated failure time (AFT) models with lognormal time distribution were used with an outcome of days to discharge alive and treatment arm as the sole covariate.²⁶ AFT-modelled associations were expressed as event time ratios (ETR) with ratios < 1 indicating shorter LOS, > 1 indicating longer LOS, and $= 1$ indicating no association of ACP with LOS. Rates of 30-day readmission by ACP status were analyzed

using Cox proportional hazard models with days to readmission as the outcome and death as a censoring event, with estimates expressed as hazard ratios.²⁷ HRs > 1 indicate a higher readmission rate, HR < 1 indicate a lower readmission rate, and HR $= 1$ indicate no association between ACP and 30-day readmission. Analysis of hospital readmission eliminated some patients due to denominator eligibility criteria and eligibility for the readmission sample could plausibly have been affected by ACP, thus separate propensity score models were fit for patients eligible for analysis of 30-day readmission (Supplemental Table 2).

Companion regression models were included with time to death as the outcome and discharge alive or 30-day readmission events as censoring events in the LOS and 30-day readmission analyses, respectively. These analyses provide context to the LOS and 30-day readmission estimates by assessing whether the relationship between ACP and LOS or 30-day readmission may have been influenced by increased or decreased in patients with ACP.

Probability of ICU transfer was analyzed using log binomial models to estimate RRs. Odds of hospice referral were modeled using multinomial logistic regression with outcome levels of discharge without referral, inpatient death, and discharge with referral.

For secondary outcomes, overlap-weighted regressions were considered primary with unweighted regressions reported for context. Percentile based CIs were calculated to account for repeated hospitalizations and uncertainty in propensity score estimation, with 1000 resamples at patient level.

Analyses used Stata Software version 17.²⁸

RESULTS

A total of 739 hospitalizations, 663 unique patients, with elevated risk of death within 30 days were extracted from the EHR; of these hospitalizations 479 met the criteria for inclusion in the analytic sample (197 encounters pre-intervention and 282 encounters post intervention) (**Figure 1**). Mean age was 75.1, 53.2% were male, and patients were predominantly White (58.2%) and Black (36.1%). A majority (59.7%) were on Medicare insurance with patient sociodemographics comparable between the pre and post period (**Table 1**). There were similar proportions of patients with medium risk scores in the pre and post periods with all “critical” patients seen in the post period.

From pre- to post-intervention periods, there was an increase in the proportion of patients receiving ACP. Adjusting for potential confounders, proportion of higher-risk patients with documented ACP rose from 6.0% (95% CI: 2.0-10.0%) at baseline to 56.5% (95% CI: 41.9, 71.1%) in the final months of the intervention (Adjusted risk difference = +50.5%, 95% CI: 36.5 – 64.6%) (**Figure 2**). Results remained similar when encounters occurring during




Figure 1. Pre and Post patient evaluation and exclusions

periods when resident teaching teams did not receive notifications were excluded from the analysis. Calculation of ICC revealed evidence of modest clustering of probability of receiving ACP by provider (ICC=0.105, 95% CI: 0.023 – 0.186).

Comparisons of standardized differences prior to weighting indicate significant imbalance in characteristics between those who received ACP and those that did not, with all standardized differences reducing to less than ± 0.10 after weighting, with the exception of provider age in the readmission sample (Supplemental Material).

Patients with ACP were more than twice as likely to have a new DNAR order placed during their admission when ACP was documented (29.0% vs. 10.8% RR=2.69, 95% CI: 1.64 – 4.27). Additionally, patients with ACP had twice the odds of hospice referral at discharge (22.2% vs. 12.6% (OR)=2.16, 95% CI: 1.16 – 4.01).

Weighted mean LOS, when discharged alive, was 7.6 days (SD=6.8) for patients with no ACP and 9.7 days (SD=9.6) for those with ACP; patients with ACP documentation had a 29% longer LOS (ETR=1.29, 95% CI: 1.10 – 1.53) compared to those without ACP. A higher weighted proportion of patients with ACP died as inpatients than those without ACP (7.2% vs. 12.4%, for no ACP and ACP, respectively; RR=1.71, 95% CI: 0.94,3.55), though time to inpatient death was comparable between those with ACP and those without (ETR=0.96, 95% CI: 0.57 – 1.51).

Proportions of patients with 30-day readmission were similar between those with ACP documentation (17.2%) and those without ACP documentation (18.5%) and

confidence intervals for all readmission related outcomes included the null ([Table 2](#)).

Provider survey

20 providers (20/41, 49%) responded to the survey. Results are presented as agree (strongly agree or agree) or disagree (strongly disagree or disagree). A majority of respondents agreed that the notifications improved care delivery (15/20, 75%), were valuable to clinical care (15/20, 75%), and that they accurately identified patients for ACP (19/20, 95%). Providers also agreed that the notification process was not difficult to navigate (17/20, 85%), was not burdensome to their daily work (13/20, 65%), is a system they would like to continue (14/20, 70%), and is something they would recommend for other specialties (18/20, 90%).

DISCUSSION

This single-center QI study evaluates the use of an ML mortality risk prediction model to improve ACP conversations and documentation for inpatient general medicine patients. It demonstrated that an ML model coupled with provider email and page notifications can increase the documentation of ACP conversations.

Similar to other studies using ML mortality risk prediction models to identify a high-risk patient population for ACP conversations for other clinical services, this project demonstrated an increase in documentation of these conversations in the EHR.^{17,18,29,30} Our study, further demonstrates the ability to utilize front-line providers to

Table 1. Patient and Provider characteristics Pre and Post intervention

	Pre-intervention	Post-intervention	Total
	(N = 197)	(N = 282)	(N = 479)
Sociodemographic Characteristics			
Age (years)			
Mean (SD)	74.0 (12.8)	75.8 (12.5)	75.1 (12.6)
Sex			
Male	106 (53.8%)	149 (52.8%)	255 (53.2%)
Race			
White	111 (56.3%)	168 (59.6%)	279 (58.2%)
Black	76 (38.6%)	97 (34.4%)	173 (36.1%)
Other race ^a	9 (4.6%)	13 (4.6%)	22 (4.6%)
Missing	1 (0.5%)	4 (1.4%)	5 (1.0%)
Marital status			
Married	80 (40.6%)	124 (44%)	204 (42.6%)
Single	31 (15.7%)	43 (15.2%)	74 (15.4%)
Divorced, separated or widowed	85 (43.1%)	108 (38.3%)	193 (40.3%)
Missing	1 (0.5%)	7 (2.5%)	8 (1.7%)
Insurance			
Private	56 (28.4%)	98 (34.8%)	154 (32.2%)
Medicaid	10 (5.1%)	13 (4.6%)	23 (4.8%)
Medicare	126 (64.0%)	160 (56.7%)	286 (59.7%)
Other/Unknown	5 (2.5%)	11 (3.9%)	16 (3.3%)
Clinical Characteristics			
Diagnosis associated with encounter			
Cardiovascular/Vascular Including Stroke	25 (12.7%)	15 (5.3%)	40 (8.4%)
Central Nervous System and/or Neurodegenerative. Non-Stroke.	14 (7.1%)	43 (15.2%)	57 (11.9%)
Fluid, Electrolyte, and hematologic abnormalities	19 (9.6%)	9 (3.2%)	28 (5.8%)
GI/Hepatology	37 (18.8%)	59 (20.9%)	96 (20.0%)
Infectious	26 (13.2%)	51 (18.1%)	77 (16.1%)
Kidney Disorders	8 (4.1%)	14 (5.0%)	22 (4.6%)
Orthopedic	7 (3.6%)	23 (8.2%)	30 (6.3%)
Other	23 (11.7%)	20 (7.1%)	43 (9.0%)
Pulmonary	38 (19.3%)	48 (17.0%)	86 (18.0%)
Risk category: 30-day risk of death			
Critical	0 (0.0%)	10 (3.5%)	10 (2.1%)
High	56 (28.4%)	66 (23.4%)	122 (25.5%)
Medium	141 (71.6%)	206 (73.0%)	347 (72.4%)
Provider Characteristics			
Age (years)			
Mean (SD)	43.4 (7.6)	42.4 (8.4)	42.8 (8.1)
Age (years)			
<35	7 (3.6%)	47 (16.7%)	54 (11.3%)
35-44	132 (67.0%)	147 (52.1%)	279 (58.2%)
45+	58 (29.4%)	88 (31.2%)	146 (30.5%)
Sex			
Male	129 (65.5%)	145 (51.4%)	274 (57.2%)
Years since residency completion			
Mean (SD)	12.4 (7.7)	10.6 (8.1)	11.3 (8.0)
Years since residency completion			
0-4 years	2 (1.0%)	52 (18.4%)	54 (11.3%)
5-12 years	111 (56.3%)	135 (47.9%)	246 (51.4%)

	Pre-intervention	Post-intervention	Total
13+ years	84 (42.6%)	95 (33.7%)	179 (37.4%)
Job title			
Assistant Professor	123 (62.4%)	122 (43.3%)	245 (51.1%)
Associate/Full Professor	24 (12.2%)	30 (10.6%)	54 (11.3%)
Medical Instructor	50 (25.4%)	130 (46.1%)	180 (37.6%)
Hospital medicine census on admission date (with interpolation)			
Mean (SD)	187.4 (11.4)	194.6 (29.2)	191.6 (23.8)
Median (Q1, Q3)	188.0 (178.0, 195.0)	197.0 (181.0, 217.0)	190.0 (180.0, 206.0)
Census per rounder			
Mean (SD)	10.6 (1.0)	10.5 (1.5)	10.5 (1.3)

^aIncludes: American Indian/Alaska Native, Asian, Other, and 2 or more races, collapsed due to sparsity




Figure 2. Regression^a adjusted^b mean proportion of high and medium risk patients receiving advanced care planning (ACP), with 95% confidence intervals

^aRegression approach is Modified Poisson with log link using generalized estimating equations clustered at the level of provider

^bEstimates are regression adjusted for: patient age, patient sex (male, female), patient marital status (single, married, divorced/separated/widowed), patient 30-day risk score (medium, high, critical) at time of admission, patient receipt of outpatient palliative care consult, disease category (cardiovascular/vascular including stroke; central nervous system and/or neurodegenerative excluding stroke, GI/hepatology, infection disease, kidney disorders, orthopedic, pulmonary, other), patient admission prior to later than 9am, provider age, provider sex (male, female), provider title (medical instructor, assistant professor, associate professor), provider years since residency completion, and hospital census per rounder on date of admission.

complete ACP conversations and documentation.³¹ This is an important finding as patients often present with acute decompensation of chronic illness or are diagnosed with a life-limiting illness during an inpatient encounter.

The substantial and sustained increase in ACP documentation in our study demonstrates how clinicians adapted ACP into their clinical workflow. Clinicians integrated the notification and ACP documentation template, where more than 50% of identified patients had ACP documented during their hospital encounter. Perhaps this is due to a manageable number of notifications clinicians received over the intervention period. Clinicians received an average of 4.3 notifications during the

analyzed intervention period. This demonstrates that with education, reminders, and notifications, frontline clinicians were able to integrate ACP documentation within their clinical workflow.³² While the model did not capture all patients who may die in six months, the model did successfully identify 238 (49%) patients who passed away in this timeframe. These notifications allowed clinicians to prioritize time to focus on ACP conversations for these high-risk patients.

Completing ACP conversations also affected code status, with an increased proportion changing from full code to DNAR prior to discharge. While a patient's code status may not communicate their goals or values, it re-

Table 2. Outcomes of encounters pre- and post- intervention

	Pre-intervention (N = 197)	Post-intervention (N = 282)	Total (N = 479)
Length of hospital stay			
Mean (SD)	8.4 (9.3)	8.9 (8.5)	8.7 (8.8)
Median (Q1, Q3)	5.9 (3.2, 9.7)	6.3 (3.8, 10.3)	6.0 (3.6, 10.1)
Transferred to ICU during hospital stay	14 (7.1%)	24 (8.5%)	38 (7.9%)
Length of stay in ICU (days)(if any ICU stay)			
Mean (SD)	5.7 (6.2)	4.2 (3.9)	4.8 (4.9)
Median (Q1, Q3)	5.2 (2.1, 6.8)	3.2 (1.0, 6.1)	3.6 (1.2, 6.8)
Inpatient death	14 (7.1%)	31 (11.0%)	45 (9.4%)
Inpatient palliative care consult during stay	38 (19.3%)	44 (15.6%)	82 (17.1%)
DNAR code status at admission	71 (36.0%)	96 (34.0%)	167 (34.9%)
DNAR	71 (36.0%)	96 (34.0%)	167 (34.9%)
Discharge disposition			
Died	14 (7.1%)	31 (11.0%)	45 (9.4%)
Hospice	32 (16.2%)	39 (13.8%)	71 (14.8%)
Home ^a	37 (18.8%)	60 (21.3%)	97 (20.3%)
Home with home care	45 (22.8%)	67 (23.8%)	112 (23.4%)
SNF, rehab, LTC, or other health facility	69 (35.0%)	85 (30.1%)	154 (32.2%)
In 30-day readmission denominator	171 (86.8%)	225 (79.8%)	396 (82.7%)

^aIncludes "Left AMA"

fects an important aspect of patient care and dictates resuscitation management in the event of cardiopulmonary arrest, illustrating one aspect of the patient's end-of-life goals.³³ We also found that patients with documented ACP were more likely to be discharged with hospice services.

Completion of ACP documentation was associated with longer LOS. This could be due to biases that we could not control for in the analysis, in which providers selected patients for ACP that they perceived to be sicker and, thus, more likely to have a longer LOS. This selection bias may be supported by the finding that patients with ACP documentation had higher inpatient and 30-day mortality in the readmission analysis.

Additionally, while our current study focuses on implementing and impacting an already developed ML model, we acknowledge the necessity of comprehensive evaluations to promote fairness across different demographic and socio-economic groups. This can be done through methodologies like transfer learning and leveraging social determinants of health data.³⁴ Though beyond the scope of this study, we highlight this as a critical direction for future research to ensure equitable benefits from ML in healthcare.

Limitations

There are several limitations to this project. First, the pre-post, non-randomized design at a single institution may limit the generalizability of the results. Future studies should incorporate randomized controlled trial designs to better evaluate clinician response to the notification.

This study used an ML model developed retrospectively on a patient cohort at a single health system, which may limit generalizability. However, the mechanism to notify hospital medicine clinicians of patients at high risk of death may be implemented in other health systems. Second, due to the pragmatic nature of this study, the measurement of ACP conversations was limited to the ascertainment of the presence of documentation, which may underestimate actual ACP conversations and does not account for the quality of these conversations. Future research should measure the quality of conversations and documentation to better evaluate the content of these conversations.²¹ Finally, though propensity score and regression adjustment models may have reduced confounding bias, our ability to make statements about causality rests on the assumption that we have included all relevant confounders, which is inherently untestable. Thus, relationships may represent associations rather than causal effects.

Conclusion

Frontline provider notifications using a ML mortality model to identify high-risk patients resulted in a sustained increase in completion of ACP documentation in the inpatient setting. This suggests that targeted provider notifications can result in increased ACP for patients with serious illness.

Author Contribution

All Authors have reviewed the final manuscript prior to submission. All the authors have contributed significantly to the manuscript, per the ICJME criteria of authorship.

- Substantial contributions to the conception or design of the work; or the acquisition, analysis, or interpretation of data for the work; AND
- Drafting the work or revising it critically for important intellectual content; AND
- Final approval of the version to be published; AND
- Agreement to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved

Funding/disclosures

Dr. Ma's effort was supported by the University of Michigan HEAL K12NS130673. The statements presented in this article are solely the responsibility of the author(s) and do not necessarily reflect the position, views, or policy of the sponsors.

Alyssa Platt's effort was supported by Grant Number UL1TR002553 from the National Center for Advancing Translational Sciences (NCATS) of the National Institutes of Health (NIH), and NIH Roadmap for Medical Research. The statements presented in this article are solely the responsibility of the author(s) and do not necessarily reflect the position, views, or policy of the sponsors.

Acknowledgement

We would like to acknowledge Chuan Hong PhD for advice and support on the statistical analysis.

Corresponding Author

Jonathan Walter MD
Department of Medicine, Division of General Internal Medicine,
Duke University School of Medicine, 40 Medicine Circle,
DUMC Box 3534, Durham, NC 27710
Phone: 270-210-2494
Email: jonathan.walter@duke.edu



This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CCBY-NC-4.0). View this license's legal deed at <https://creativecommons.org/licenses/by-nc/4.0> and legal code at <https://creativecommons.org/licenses/by-nc/4.0/legalcode> for more information.

REFERENCES

1. Sudore RL, Lum HD, You JJ, et al. Defining Advance Care Planning for Adults: A Consensus Definition From a Multidisciplinary Delphi Panel. *J Pain Symptom Manage*. 2017;53(5):821-832.e1. doi:10.1016/j.jpainsymman.2016.12.331
2. Wright AA, Zhang B, Ray A, et al. Associations between end-of-life discussions, patient mental health, medical care near death, and caregiver bereavement adjustment. *JAMA*. 2008;300(14):1665-1673. doi:10.1001/jama.300.14.1665
3. Bharadwaj P, Helfen KM, Deleon LJ, et al. Making the Case for Palliative Care at the System Level: Outcomes Data. *Journal of Palliative Medicine*. 2016;19:255-258. doi:10.1089/jpm.2015.0234
4. May P, Normand C, Cassel JB, et al. Economics of Palliative Care for Hospitalized Adults With Serious Illness: A Meta-analysis. *JAMA Intern Med*. 2018;178:820-829. doi:10.1001/jamainternmed.2018.0750
5. Morrison RS, Penrod JD, Cassel JB, et al. Cost savings associated with US hospital palliative care consultation programs. *Archives of Internal Medicine*. 2008;168:1783-1790. doi:10.1001/archinte.168.16.1783
6. O'Connor NR, Junker P, Appel SM, Stetson RL, Rohrbach J, Meghani SH. Palliative Care Consultation for Goals of Care and Future Acute Care Costs: A Propensity-Matched Study. *Am J Hosp Palliat Care*. 2018;35(7):966-971. doi:10.1177/1049909117743475
7. Yadav KN, Gabler NB, Cooney E, et al. Approximately One In Three US Adults Completes Any Type Of Advance Directive For End-Of-Life Care. *Health Aff (Millwood)*. 2017;36(7):1244-1251. doi:10.1377/hlthaff.2017.0175
8. Sadeghi B, Walling AM, Romano PS, Ahluwalia SC, Ong MK. A Hospital-Based Advance Care Planning Intervention for Patients with Heart Failure: A Feasibility Study. *J Palliat Med*. 2016;19(4):451-455. doi:10.1089/jpm.2015.0269
9. Lamas D, Panariello N, Henrich N, et al. Advance Care Planning Documentation in Electronic Health Records: Current Challenges and Recommendations for Change. *J Palliat Med*. 2018;21(4):522-528. doi:10.1089/jpm.2017.0451
10. You JJ, Downar J, Fowler RA, et al. Barriers to Goals of Care Discussions With Seriously Ill Hospitalized Patients and Their Families: A Multicenter Survey of Clinicians. *JAMA Intern Med*. 2015;175(4):549-556. doi:10.1001/jamainternmed.2014.7732
11. Howard M, Bernard C, Klein D, et al. Barriers to and enablers of advance care planning with patients in primary care: Survey of health care providers. *Can Fam Physician*. 2018;64(4):e190-e198.
12. Ouchi K, Strout T, Haydar S, et al. Association of Emergency Clinicians' Assessment of Mortality Risk With Actual 1-Month Mortality Among Older Adults Admitted to the Hospital. *JAMA Netw Open*. 2019;2(9):e1911139. doi:10.1001/jamanetworkopen.2019.11139
13. Christakis NA, Lamont EB. Extent and determinants of error in doctors' prognoses in terminally ill patients: prospective cohort study. *BMJ*. 2000;320(7233):469-472. doi:10.1136/bmj.320.7233.469
14. Parikh RB, Kakad M, Bates DW. Integrating Predictive Analytics Into High-Value Care: The Dawn of Precision Delivery. *JAMA*. 2016;315(7):651-652. doi:10.1001/jama.2015.19417
15. Parikh RB, Manz C, Chivers C, et al. Machine Learning Approaches to Predict 6-Month Mortality Among Patients With Cancer. *JAMA Netw Open*. 2019;2(10):e1915997. doi:10.1001/jamanetworkopen.2019.15997
16. Wang E, Major VJ, Adler N, et al. Supporting Acute Advance Care Planning with Precise, Timely Mortality Risk Predictions. *NEJM Catalyst Innovations in Care Delivery*. 2021;03. doi:10.1056/CAT.20.0655
17. Manz CR, Parikh RB, Small DS, et al. Effect of Integrating Machine Learning Mortality Estimates With Behavioral Nudges to Clinicians on Serious Illness Conversations Among Patients With Cancer: A Stepped-Wedge Cluster Randomized Clinical Trial. *JAMA Oncol*. 2020;6(12):e204759. doi:10.1001/jamaoncol.2020.4759
18. Courtright KR, Chivers C, Becker M, et al. Electronic Health Record Mortality Prediction Model for Targeted Palliative Care Among Hospitalized Medical Patients: A Pilot Quasi-experimental Study. *J Gen Intern Med*. 2019;34(9):1841-1847. doi:10.1007/s11606-019-05169-2
19. Brajer N, Cozzi B, Gao M, et al. Prospective and External Evaluation of a Machine Learning Model to Predict In-Hospital Mortality of Adults at Time of Admission. *JAMA Network Open*. 2020;3(2):e1920733. doi:10.1001/jamanetworkopen.2019.20733
20. DIHI Mortality Model Information/Evaluation. Duke Institute for Health Innovation. Accessed May 12, 2023. https://dihi.org/model-fact-sheets/docs_mortality_evaluation
21. Schlichte L, Setji N, Walter J, et al. The Use of Templates for Documenting Advance Care Planning Conversations: A Descriptive Analysis. *J Pain Symptom Manage*. 2023;66(2):123-136. doi:10.1016/j.jpainsymman.2023.04.015

22. Zou GY, Donner A. Extension of the modified Poisson regression model to prospective studies with correlated binary data. *Stat Methods Med Res.* 2013;22(6):661-670. [doi:10.1177/0962280211427759](https://doi.org/10.1177/0962280211427759)
23. Wu S, Crespi CM, Wong WK. Comparison of methods for estimating the intraclass correlation coefficient for binary responses in cancer prevention cluster randomized trials. *Contemp Clin Trials.* 2012;33(5):869-880. [doi:10.1016/j.cct.2012.05.004](https://doi.org/10.1016/j.cct.2012.05.004)
24. Li F, Thomas LE, Li F. Addressing extreme propensity scores via the overlap weights. *Am J Epidemiol.* 2019;188(1):250-257. [doi:10.1093/aje/kwy201](https://doi.org/10.1093/aje/kwy201)
25. Yang D, Dalton JE. A unified approach to measuring the effect size between two groups using SAS. *SAS global forum.* 2012;335(1):1.
26. Cho Y, Hu C, Ghosh D. Covariate adjustment using propensity scores for dependent censoring problems in the accelerated failure time model. *Stat Med.* 2018;37(3):390-404. [doi:10.1002/sim.7513](https://doi.org/10.1002/sim.7513)
27. Cheng C, Li F, Thomas LE, Li FF. Addressing Extreme Propensity Scores in Estimating Counterfactual Survival Functions via the Overlap Weights. *Am J Epidemiol.* 2022;191(6):1140-1151. [doi:10.1093/aje/kwac043](https://doi.org/10.1093/aje/kwac043)
28. *Stata Software.* Stata Corporation
29. Manz CR, Zhang Y, Chen K, et al. Long-term Effect of Machine Learning–Triggered Behavioral Nudges on Serious Illness Conversations and End-of-Life Outcomes Among Patients With Cancer: A Randomized Clinical Trial. *JAMA Oncol.* 2023;9(3):414-418. [doi:10.1001/jamaoncol.2022.6303](https://doi.org/10.1001/jamaoncol.2022.6303)
30. Schell JO, Schenker Y, Piscitello G, et al. Implementing a Serious Illness Risk Prediction Model: Impact on Goals of Care Documentation. *J Pain Symptom Manage.* 2023;66(6):603-610.e3. [doi:10.1016/j.jpainsymman.2023.07.015](https://doi.org/10.1016/j.jpainsymman.2023.07.015)
31. Chi S, Kim S, Reuter M, et al. Advanced Care Planning for Hospitalized Patients Following Clinician Notification of Patient Mortality by a Machine Learning Algorithm. *JAMA Netw Open.* 2023;6(4):e238795. [doi:10.1001/jamanetworkopen.2023.8795](https://doi.org/10.1001/jamanetworkopen.2023.8795)
32. Ma JE, Haverfield M, Lorenz KA, et al. Exploring expanded interdisciplinary roles in goals of care conversations in a national goals of care initiative: A qualitative approach. *Palliat Med.* 2021;35(8):1542-1552. [doi:10.1177/02692163211020473](https://doi.org/10.1177/02692163211020473)
33. Bhatia HL, Patel NR, Choma NN, Grande J, Giuse DA, Lehmann CU. Code status and resuscitation options in the electronic health record. *Resuscitation.* 2015;87(33):14-20. [doi:10.1016/j.resuscitation.2014.10.022](https://doi.org/10.1016/j.resuscitation.2014.10.022)
34. Gao Y, Cui Y. Deep transfer learning for reducing health care disparities arising from biomedical data inequality. *Nat Commun.* 2020;11(1):5131. [doi:10.1038/s41467-020-18918-3](https://doi.org/10.1038/s41467-020-18918-3)

SUPPLEMENTARY MATERIALS

Supplementary Material

Download: https://bhm.scholasticahq.com/article/120907-quality-improvement-study-using-a-machine-learning-mortality-risk-prediction-model-notification-system-on-advance-care-planning-in-high-risk-patients/attachment/235426.docx?auth_token=tXEM3DiPfyUc1WpWHEID
